



## Endocrine complications of celiac disease: a case report and review of the literature

Marcella D. Walker, Haley M. Zylberberg, Peter H. R. Green & Michael S. Katz

To cite this article: Marcella D. Walker, Haley M. Zylberberg, Peter H. R. Green & Michael S. Katz (2018): Endocrine complications of celiac disease: a case report and review of the literature, Endocrine Research, DOI: [10.1080/07435800.2018.1509868](https://doi.org/10.1080/07435800.2018.1509868)

To link to this article: <https://doi.org/10.1080/07435800.2018.1509868>



Published online: 10 Sep 2018.



Submit your article to this journal [↗](#)



Article views: 342



View Crossmark data [↗](#)



## Endocrine complications of celiac disease: a case report and review of the literature

Marcella D. Walker<sup>a</sup>, Haley M. Zylberberg<sup>b</sup>, Peter H. R. Green<sup>a</sup>, and Michael S. Katz<sup>c</sup>

<sup>a</sup>Department of Medicine, Columbia University, New York, NY, USA; <sup>b</sup>Department of Medicine, Mount Sinai, New York, NY, USA; <sup>c</sup>Department of Medicine, University of Texas Health Science Center at San Antonio, San Antonio, TX, USA

### ABSTRACT

**Purpose:** The purpose of this article is to review recent literature regarding endocrine disorders related to celiac disease (CD).

**Methods:** We describe a case report and review existing literature on the endocrine manifestations of CD.

**Results:** CD is an autoimmune disorder characterized by intestinal inflammation in response to gluten. CD can cause a wide range of extra-intestinal complications, including endocrine manifestations. Metabolic bone disease including osteoporosis and osteopenia, vitamin D deficiency, secondary hyperparathyroidism and less frequently osteomalacia can be seen. In CD, fracture risk is increased by 30–40%, while risk for hip fracture is approximately doubled. The risk for other endocrine disorders, particularly autoimmune endocrinopathies, is also increased in those with CD compared to the general population. Epidemiologic data indicate the risk for hypothyroidism is 3–4 times higher among those with CD, while risk of type 1 diabetes is greater than double. Risk for primary adrenal insufficiency is a striking 11-fold higher in those with versus without CD, though the absolute risk is low. Fertility is reduced in women with CD before diagnosis by 37% while male fertility in the absence of hypogonadism does not appear to be affected. Other endocrine conditions including hyperthyroidism, ovarian failure, androgen insensitivity, impaired growth and growth hormone deficiency and autoimmune polyendocrine syndromes have also been associated with CD.

**Conclusions:** CD is associated with a wide range of endocrine manifestations.

### ARTICLE HISTORY

Received 20 June 2018

Revised 31 July 2018

Accepted 4 August 2018

### KEYWORDS

Celiac disease; endocrine manifestations; hypothyroidism; osteoporosis

### Introduction

CD is an autoimmune disorder characterized by intestinal inflammation in response to gluten, a family of proteins found in wheat, barley, and rye.<sup>1</sup> Classic CD is characterized by malabsorption, diarrhea, and weight loss, though today many patients have only minor gastrointestinal symptoms and can be asymptomatic. CD can also cause a wide range of extra-intestinal complications, including endocrine manifestations.<sup>2</sup> Metabolic bone disease including osteoporosis and osteopenia, vitamin D deficiency, secondary hyperparathyroidism and less frequently osteomalacia can be seen. CD is also associated with other endocrine disorders including, hypo- and hyperthyroidism, type 1 diabetes, adrenal insufficiency, ovarian failure and infertility, androgen insensitivity, impaired growth and growth hormone deficiency (GHD) and autoimmune polyendocrine syndromes. This review will

summarize endocrine disorders related to CD with a focus on recent investigations where available.

### Methods

We describe a case report of a patient with CD and multiple endocrine comorbidities and review the existing literature of the endocrine manifestations of CD.

### Case report

A 60-year-old man presented for endocrine evaluation due to a 4-year history of fatigue, reduced muscle mass and strength, loss of libido, mild cognitive changes, depressive symptoms, and weight loss of 20 pounds. Prior initial outside evaluation indicated an elevated total testosterone level. The patient reported loss of appetite and constipation but no vomiting, diarrhea, abdominal pain, or food

intolerance. Colonoscopy, 3 years prior, had been normal. He was the biological father of two sons ages 30 and 27 and reported no other symptoms of hypogonadism. He described one inch of height loss without history of fracture and did not smoke or consume alcohol. He had been on varying antiepileptic drugs for 30 years following a generalized tonic-clonic seizure associated with a right temporal subarachnoid cyst, which was stable on serial imaging. At presentation, his medications included

divalproex and vitamin D. The patient denied the use of other medications.

On examination, the patient was 67.5 kilograms and euthyroid-appearing with a normal adult male phenotype and body mass index (BMI) 22.1 kg/m<sup>2</sup>. He did not meet criteria for anorexia. The thyroid gland was normal in size with irregular texture. There was no kyphosis or gynecomastia. Abdominal exam was normal. Circumcised phallus and testes were normal, the latter measuring 4.5 × 3.0 cm each.

**Table 1.** Case report laboratory evaluation.

	Normal Range	Initial Evaluation Pre-treatment	1-year after Testosterone Treatment	After GFD Initiation on Testosterone	On GFD and off Testosterone
<b>Serum Biochemistries, Renal function, Liver function</b>					
Sodium (mmol/L)	135–145	132, 138			
Potassium (mmol/L)	3.3–5.0	4.2			
Creatinine (mg/dL)	0.5–1.4	1.3, 1.1	1.0		
Glucose (mg/dL)	64–99	88			
Albumin (g/dL)	3.1–4.7	4.0	4.7		4.8
AST (U/L)	3–70	38			
ALT (U/L)	3–78	38			
Alkaline Phosphatase (U/L)	20–150	80	220		69
<b>Blood count and Iron Studies</b>					
Hematocrit (%)	42.0–53.0	40.7, 43.0			
Hemoglobin (gm/dL)	14.5–17.7	14.0, 14.8			
Serum Iron (µg/dL)	49–181	97			
TIBC (µg/ml)	228–428	344			
Iron Saturation (%)	20–55	28			
Vitamin B12 (pg/mL)	211–911	492			
<b>Metabolic Bone Evaluation</b>					
Serum Calcium (mg/dL)	8.4–10.5	9.1	9.5		9.7
Ionized Calcium (mg/dL)	4.6–5.4		4.8		4.8
Serum Phosphorus (mg/dL)	2.4–5.0		4.0		
25-hydroxyvitamin D (ng/mL)	20–50		35		55
1,25-dihydroxyvitamin D	18–64		78		
PTH (pg/mL)	14–72		155		47.5
24-hour urine calcium (mg/24 hours)	100–250		< 26		80–164
Bone-specific Alkaline Phosphatase (mcg/L)	0–20		35		
NTX (nmol BCE/mmol creatinine)	21–66		209	32	
<b>Hormones and Endocrine Testing</b>					
Cortisol (µg/dL)	4.30–22.40	11.04			
TSH (µIU/mL)	0.40–5.00	1.00	1.13		
TPO antibodies (IU/mL)	< 9.0			44.5	15.7
Total Testosterone (ng/dL)	240–950	1030, 1120	1280, 953	958	831
Free Testosterone (ng/dL)	9–30	15.5, 9.0	17, 13	15	10.8
SHBG (nmol/L)	10–60	145, 160	116	83	84
LH (mIU/mL)	1.5–9.3	37.3, 46.6	16.8	5.6	22.8
FSH (mIU/mL)	1.4–18.1	19.8, 22.9	13.6	8.1	22.3
Prolactin (ng/mL)	2.1–17.7	3.8			
Estradiol (pg/mL)	11.6–41.2	< 11.8			
Inhibin B (pg/mL)	< 399	320			
<b>Celiac Serologies</b>					
tTG IgA (U/L)	< 4.0		5.4		
Deaminated gliadin IgG (U)	< 20		19.6		
Total IgA(mg/dL)	61–356		9		

GFD = gluten-free diet; TIBC = total iron binding capacity; PTH = parathyroid hormone; NTX = n-telo-peptide; TSH = thyroid-stimulating hormone; SHBG = sex hormone binding globulin; LH = lutenizing hormone; FSH = follicle-stimulating hormone; tTG = transglutaminase

On initial laboratory testing (Table 1), serum glucose, renal, liver and thyroid function, electrolytes, calcium, albumin, alkaline phosphatase, and morning cortisol levels were all normal. Hemoglobin and hematocrit were borderline low to frankly low with normal vitamin B12 level and iron studies, which were not consistent with hemochromatosis. As shown in Table 1, initial and repeat endocrine testing (Mayo Clinic Laboratories, Rochester, MN) revealed total testosterone ranging from 1030 to 1120 ng/dL (adult male reference range: 240–950 ng/dL), free testosterone 9.0–15.5 ng/dL (ref: 9–30), luteinizing hormone (LH) 37.3–46.6 mIU/mL (ref: 1.5–9.3), follicle stimulating hormone (FSH) 19.8–22.9 mIU/mL (ref: 1.4–18.1), sex hormone-binding globulin (SHBG) 145–160 nmol/L (ref: 10–60), prolactin 3.8 ng/mL (ref: 2.1–17.7), estradiol <11.8 pg/mL (ref: 11.6–41.2), and inhibin B 320 pg/mL (ref: <399).

As the patient's symptoms were consistent with hypogonadism, free testosterone levels were generally in the lower reference range (despite elevated total testosterone with high SHBG levels) and gonadotropins were elevated, a trial of transdermal testosterone was initiated. Over one year, testosterone supplementation with titration had no effect on the patient's symptoms, weight or total or free testosterone (Table 1); in contrast, LH and FSH fell to 16.8 and 13.6 mIU/mL, respectively, with SHBG 116 nmol/L (following a change of divalproex to phenytoin).

Dual energy x-ray absorptiometry (DXA; Lunar Prodigy, GE Healthcare, Madison, WI) performed one year after testosterone initiation due to his history of anti-epileptic drug use revealed osteoporosis with T-scores of -5.2, -3.8, -3.4, and -4.3 at the lumbar spine, hip, femoral neck and 1/3 radius, respectively. Z-scores were below -2.0 at all skeletal sites (-4.8, -3.3, -2.4, and -3.8 at the lumbar spine, hip, femoral neck and 1/3 radius, respectively). Spinal X-rays were not performed. Due to extremely low T-scores, an evaluation for secondary causes of osteoporosis was performed (Table 1). Serum total (9.5 mg/dL; ref: 8.4–10.5) and ionized calcium (4.8 mg/dL; ref: 4.6–5.4) levels were normal as was phosphorus (4.0 mg/dL; ref: 2.4–5.0), renal function, albumin (4.7 g/dL; ref: 3.1–4.7), and 25-hydroxyvitamin D (35 ng/mL) on supplementation. In contrast, 1,25(OH)<sub>2</sub> vitamin D (78 pg/mL; ref: 18–64) and intact

parathyroid hormone (iPTH) 155 pg/mL (ref: 14–72) were elevated. Calcium excretion based on a 24-hour urine collection (total volume 2550 ml and creatinine 1.4 g) was undetectable (<26 mg; ref: 100–250). Total alkaline phosphatase (220 U/L; ref: 20–150), bone-specific alkaline phosphatase (35 mcg/L; ref: 0–20) and urine n-telopeptide (NTX; 209 nmol BCE/mmol creatinine; ref: 21–66) were elevated. HIV testing was not performed. Thyroid function tests were again normal. The laboratory findings were interpreted as indicating secondary hyperparathyroidism due to calcium malabsorption complicated by osteomalacia.

Celiac disease (CD) screening included an evaluation for secondary causes of bone loss revealed the following: CD serologies were equivocally positive in the setting of IgA deficiency (total IgA 9 mg/dL; ref: 61–356) with tissue transglutaminase (tTG) IgA 5.4 U/mL (<4.0 negative) and deamidated gliadin IgG 19.6 U (<20.0 negative); human leukocyte antigen (HLA) testing was positive for DQ2 alleles associated with CD. A duodenal biopsy by endoscopy showing severe villous atrophy and lymphocytosis confirmed CD. The diagnosis of CD with metabolic bone disease and androgen resistance was made 2 years after referral to endocrinology and six years after symptom onset.

A gluten-free diet (GFD) was initiated and rigorously maintained. Successive biopsies revealed duodenal mucosal healing by 14 months on GFD. Constitutional symptoms and constipation improved and the patient gained 20 pounds after 3 years on GFD. Bone mineral density (BMD) increased over 2 years by 36.5%, 21.8%, and 14.2% at the spine, hip, and 1/3 radius, respectively, while on a GFD with supplemental calcium and vitamin D. Within about the same period, metabolic bone indices normalized (Table 1): iPTH 47.5 pg/mL, alkaline phosphatase 69 U/L, 25OH vitamin D 55 ng/mL, 24-hour urine calcium 80–164 mg/24 hrs, and urine NTX 32 nmol BCE/mmol creatinine. During the first 16 months of GFD and on tapering testosterone supplementation, both LH and FSH normalized (5.6 and 8.1 mIU/mL, respectively), while SHBG decreased to 83 nmol/L without a change in total and free testosterone.

Subsequently, on continued GFD without testosterone, LH and FSH rose (22.8 and 22.3 mIU/mL, respectively) without substantial changes in total or free testosterone or SHBG (Table 1). Notwithstanding multiple confounders, the serial changes in gonadotropins and SHBG were interpreted as incomplete androgen insensitivity partially alleviated by GFD. On GFD, the patient remained euthyroid with mildly elevated thyroid peroxidase antibodies up to 44.5 IU/mL (ref: <9.0) and stable, bilateral sub-centimeter thyroid nodules by ultrasound. Due to T-scores remaining in the osteoporotic range, the patient was treated with teriparatide followed by zoledronic acid.

### Epidemiology, pathogenesis, and diagnosis

The global prevalence of CD has historically been estimated to be 1%, based mostly on European and other Caucasian populations.<sup>1,3</sup> A recent meta-analysis that included 96 studies and 275,818 individuals suggests CD has a pooled worldwide prevalence of 1.4% and 0.7% based on serologies and biopsy data, respectively;<sup>4</sup> the prevalence, however, varies geographically and ranges from 1.1% in Africa (based on serology) to 1.8% in Asia with intermediate values in Europe, the Americas, and Oceania.<sup>4</sup> Within the United States, CD is more common in Caucasians of European ancestry and less frequent among African-Americans and Hispanic-Americans.<sup>5</sup> Even within individual European countries, the prevalence can vary considerably.<sup>3</sup> Recent work suggests CD is common in northern India and newer data suggests it may not be uncommon in Southeast Asia.<sup>4,6</sup> Both in the United States and Western Europe the prevalence of CD has increased since the 1950s, likely due to a combination of increased global incidence, increased patient and physician awareness, and increases in CD diagnosis.<sup>1,7-9</sup>

CD can be diagnosed at any age, but recent work suggests the prevalence is almost twice as high in children compared to adults.<sup>4,9</sup> There is a female predominance of the disease, though this may be due to disparities in diagnosis as opposed to an inherent feature of the disease.<sup>4,10</sup> CD has a strong genetic basis – 90% of individuals with CD carry a particular HLA genotype, HLA-DQ2, and the remainder carry HLA-DQ8.<sup>1,11</sup> Nevertheless, not all individuals who are positive for HLA-DQ2 or HLA-DQ8 develop CD

upon exposure to gluten, and so genes outside of the HLA loci and non-dietary environmental factors are thought to contribute to the pathogenesis.<sup>1,11</sup> Recent investigations point to the possible role of the gut microbiome in contributing to development of CD.<sup>12</sup>

Individuals with CD develop an immune response to the ingestion of gluten, specifically its peptide component gliadin. In the intestinal lamina propria, the enzyme tissue transglutaminase (tTG) deamidates gliadin, enabling its binding to HLA-DQ2 or HLA-DQ8 molecules on antigen presenting cells and presentation to CD4 + T cells.<sup>11</sup> Proinflammatory cytokines and autoantibodies are subsequently produced in this adaptive immune response. The innate immune system is also involved and is reflected in the increase in intraepithelial lymphocytes, which express natural killer T lymphocyte receptors.<sup>11</sup> It is unclear how the adaptive lamina propria immune response and the intraepithelial innate immune reactions relate to each other, but it is apparent that all are required for the “full blown” intestinal celiac lesion characterized by villous atrophy and intraepithelial lymphocytosis.<sup>1,2,11</sup>

Diagnosis of CD relies on a combination of testing for elevated concentrations of autoantibodies and duodenal biopsy showing villous atrophy and intraepithelial lymphocytosis.<sup>1</sup> Autoantibodies used in screening include IgA-tTG, which has a high sensitivity and negative predictive value, as well as IgA endomysial antibodies (EMA) and IgA and IgG deamidated gliadin peptides.<sup>1,13</sup> In adults, it is generally recommended that serologic testing for CD be performed first, and then if positive, the diagnosis should be confirmed by duodenal biopsy.<sup>1</sup> In the pediatric population, recent European guidelines allow for serological diagnosis without biopsy confirmation in some cases.<sup>14</sup> Following diagnosis, patients with CD are instructed to adhere to a life-long gluten-free diet (GFD), which is the only currently available treatment.

### Metabolic bone disease

Metabolic bone disease is reported to be a frequent complication of CD and is sometimes the presenting manifestation. Multiple reports suggest that osteopenia and osteoporosis are common in CD and screening with DXA is recommended at diagnosis.<sup>15</sup> There is, however, a wide range of

prevalence estimates described, varying from 9 to 75%.<sup>16–19</sup> Further, a recent meta-analysis suggests that the rate of CD among those with osteoporosis is 1 in 62 or about 1.6%, which is comparable to that in the general population.<sup>20</sup> Low BMD can occur in both those with and without gastrointestinal symptoms, such as diarrhea.<sup>17,21,22</sup> Data regarding fracture risk in patients with CD is conflicting but a recent meta-analysis of prospective studies indicated that CD was associated with a 30% increase in the risk of any fracture and a 69% increase in the risk (Table 2) of hip fracture.<sup>23,24,33–36</sup> Another recent study suggests CD is associated with an increased fracture risk assessment (FRAX) score, implying it is an independent risk factor for fracture.<sup>22</sup>

Recent advances in skeletal imaging, including the development of high resolution peripheral quantitative computed tomography (HRpQCT), have allowed insight into the pathogenesis of skeletal fragility in CD. A recent HRpQCT study in adult women with CD indicated low trabecular volumetric BMD and fewer, more widely and irregularly spaced trabeculae at both the radius and tibia as well as thinner cortices at the tibia versus controls.<sup>37</sup> These microstructural deficits led to lower whole-bone stiffness and failure load in CD patients at both skeletal sites, which would

predispose to an increased risk of fracture. A similar study in premenopausal women likewise found trabecular deficits and lower cortical density at both skeletal sites but no difference in cortical thickness.<sup>38</sup> The latter study also suggested that those with symptomatic versus asymptomatic CD had greater microstructural deterioration.<sup>38</sup>

The pathogenesis of reduced BMD, microarchitectural deterioration, and increased skeletal fragility in CD remains incompletely investigated, but both local and systemic mechanisms have been implicated.<sup>15,39</sup> The clinical features of CD-associated metabolic bone disease are shown in Table 3. While metabolic bone disease is one thought to be one of the most frequent complications of CD, few studies have directly investigated the mechanism and more research is needed in this area. Weight loss, general malnutrition, and hypogonadism may contribute to bone loss in some CD patients. A major mechanism, however, is thought to be villous atrophy in the small bowel, the major site of calcium and vitamin D absorption. In some cases, malabsorption of calcium and vitamin D may lead to compensatory secondary hyperparathyroidism and bone resorption to maintain normal serum calcium levels.<sup>15,21,39</sup> Although there are limited data regarding the prevalence of secondary hyperparathyroidism in CD, one study indicates it is present in 28% of patients and that PTH level is negatively associated with BMD at the forearm.<sup>40</sup> A second study suggested a prevalence of 25% in newly diagnosed patients as well as those refractory to treatment while the rate was 19% in those who were responsive to treatment.<sup>41</sup> In some cases, prolonged secondary hyperparathyroidism can evolve into autonomous parathyroid function and primary hyperparathyroidism (PHPT). A recent study suggests that CD patients are at almost twice the risk for PHPT.<sup>42</sup> Osteomalacia may be present in some patients due to prolonged vitamin D deficiency and calcium malabsorption, though it appears to be rare today.<sup>43</sup>

Another contributor to bone loss in CD may be the underlying autoimmune and inflammatory component of CD. CD increases serum levels of proinflammatory cytokines interleukin-1 and -6 and tumor necrosis factor-alpha, which could increase the ratio of receptor activator of nuclear factor kappa B ligand (RANKL) to osteoprotegerin (OPG).<sup>15,44</sup> High RANKL/OPG ratios increase the

**Table 2.** Risk of endocrine disease in celiac disease.

Endocrine Manifestation	Risk Estimates for Various Endocrine Diseases in CD vs. the general population
Fracture	
Any type	HR 1.4 (95% CI 1.3–1.5) <sup>23</sup> HR 1.3 (95% CI 1.14–1.5) <sup>24</sup>
Hip Fracture	HR 2.1 (95% CI 1.8–2.4) <sup>23</sup> HR 1.69 (95% CI 1.1–2.59) <sup>24</sup>
Thyroid Disease	
Any type of thyroid disease	OR 3.08 (95% CI 2.67–3.56) <sup>25</sup>
Hypothyroidism	OR 3.38 (95% CI 2.88–6.56) <sup>25</sup> HR 4.4 (95% CI 3.4–5.6) <sup>26</sup>
Hyperthyroidism	OR 1.28 (95% CI 0.37–4.46) <sup>25</sup> HR 2.9 (95% CI 2.0–4.2) <sup>26</sup>
Euthyroid autoimmune thyroid disease (i.e. + Thyroid antibodies)	OR 4.35 (95% CI 2.88–6.56) <sup>25</sup>
Type 1 Diabetes	HR 2.4 (95% CI 1.9–3.0) <sup>27</sup>
Addison's Disease	HR 11.4 (4.4–29.6) <sup>28</sup>
Female Fertility	
Overall (Pre- and Post-diagnosis CD)	RR 1.01 (95% CI 0.91–1.14) <sup>29</sup> HR 1.03 (95% CI 1.01–1.05) <sup>30</sup>
Pre-diagnosis CD	HR = 0.63 (95% CI 0.57–0.70) <sup>30</sup>
Male Infertility	HR 1.02 (0.99–1.04) <sup>31</sup> OR 1.63 (0.66–4.46) <sup>32</sup>

OR = Odds Ratio; HR = Hazard Ratio; CI = Confidence Interval; CD = Celiac Disease; RR = Rate Ratio

**Table 3.** Major features of endocrine conditions associated with celiac disease.

Endocrine Comorbidity	Possible Clinical Features
Celiac-Associated Bone Disease	Osteopenia or osteoporosis by DXA; increased risk of fracture; vitamin D deficiency or insufficiency; in some cases secondary hyperparathyroidism is present; rarely osteomalacia
Hypothyroidism	Weight gain; lethargy/fatigue, low mood; bradycardia; cold intolerance; constipation; dry skin; hair loss; irregular menses; slow growth in children; goiter; facial swelling; hoarseness
Hyperthyroidism	Weight loss; tachycardia; fever; heat intolerance; myopathy; hyperactivity/restlessness; anxiety; tremulousness; insomnia; diarrhea; hair loss; smooth, fine skin; irregular menses; goiter
Euthyroid Hashimoto's Thyroiditis	Positive TPO antibodies; normal thyroid function tests; heterogeneous echotexture of thyroid on ultrasound; increased risk for progression to hypothyroidism;
Type 1 Diabetes	Weight loss; increased thirst; increased urination; hunger; fatigue; blurred vision; nausea; vomiting; pre-syncope/dehydration
Addison's Disease	Weight loss; skin hyperpigmentation; fatigue; anorexia; abdominal pain; nausea; vomiting; hypotension/shock; presyncope; dehydration; hypoglycemia; electrolyte abnormalities including hyponatremia and hyperkalemia
Hypogonadism	
Female	Irregular menses; amenorrhea; infertility
Male	Decreased energy and libido; erectile dysfunction; decreased muscle mass; reduced terminal hair growth; low bone mineral density; infertility
Androgen resistance	Clinical features of male hypogonadism; increased total testosterone, SHBG and LH; low DHT
Growth Hormone Deficiency	
Adult	Low energy/quality of life; decreased muscle strength and exercise tolerance; increased weight/fat mass; reduced muscle mass; low bone mass
Childhood	Delayed growth/short stature; weight:height ratios increased; infantile fat distribution; delayed bone age
Autoimmune Polyendocrine Syndrome 2	Features of Addison's disease, Type 1 diabetes and thyroid disease as described; alopecia, vitiligo, pernicious anemia and primary ovarian failure may be associated; auto-antibodies against thyroid peroxidase, glutamic acid decarboxylase 65, and to 21-hydroxylase may be positive

DXA = dual x-ray absorptiometry, TPO = thyroid peroxidase, SHBG = sex hormone binding globulin, LH = luteinizing hormone; DHT = dihydrotestosterone

maturation and activation of osteoclasts, which are responsible for bone resorption. There are limited data regarding whether OPG or RANKL levels are altered in patients with CD, though one report indicates that the OPG/RANKL ratio is lower in CD patients than in controls and was positively correlated with spine BMD.<sup>45</sup> Another study demonstrated OPG neutralizing antibodies are present in some patients with CD.<sup>46</sup>

With institution of and adherence to a GFD, BMD tends to improve in CD patients, with much of the recovery taking place in the first year.<sup>47–56</sup> There is considerable intra-individual variability, however, and little data about which patients have the greatest capacity for skeletal recovery. Some data suggest improvement only in symptomatic CD while another study showed similar degrees of BMD recovery in those without symptoms.<sup>21,49</sup> Other studies suggest that improvement depends on menopausal status or age.<sup>49,53,54,57</sup> A recent study in a large celiac cohort suggested that those with the lowest serum calcium had the greatest BMD increase after diagnosis consistent with calcium malabsorption influencing the

potential for recovery.<sup>43</sup> Despite improvement, however, GFD adherence may not completely restore BMD and increased fracture risk persists even after institution of a GFD.<sup>17,23,36,48,52</sup> Recent data suggest those with persistent villous atrophy remain at increased risk after treatment compared to those with mucosal healing.<sup>34,39,58</sup> A 2017 study found that more severe Marsh histopathological stage, a system of measuring severity of intestinal damage in CD, was associated with low BMD.<sup>59</sup>

There are almost no data assessing the effect of pharmacological osteoporosis treatment specifically in patients with CD.<sup>60</sup> In the absence of data, it is reasonable to consider treatment in postmenopausal women and men >age 50 with osteoporosis that does not improve on a GFD, those with a fragility fracture or those at high risk of fracture, based on National Osteoporosis Foundation (NOF) guidelines. Parenteral agents may theoretically be more appealing due to the possibility of inadequate absorption of bisphosphonates in the setting of CD. However, anti-resorptives are contraindicated in those with hypocalcemia, vitamin D deficiency,

or osteomalacia and can precipitate or worsen hypocalcemia.<sup>61</sup> Mineral metabolism abnormalities, including secondary hyperparathyroidism, should be corrected prior to initiation of therapy.

## Association with other endocrine diseases

### Thyroid disease

Both hypo- and hyperthyroidism have been associated with CD (Table 2). The clinical features of thyroid disease are shown in Table 3. The link between autoimmune thyroid disease (AIT) and CD is incompletely understood but is likely due in part to HLA genotypes predisposing to both conditions. HLA-DQ2 and DQ8 are associated with both CD and AIT.<sup>62</sup> Non-HLA loci have been implicated as well. Variants in the gene encoding cytotoxic T-lymphocyte-associated antigen-4 (CTLA-4), which serves as an immune checkpoint and downregulates immune responses, have also been linked to both CD and AIT disease.<sup>63–65</sup> Inconsistent data indicate adherence to a GFD may reduce the risk of developing additional autoimmune disorders suggesting that uncontrolled chronic inflammation could also play a role in promoting more generalized endocrine autoimmunity.<sup>66,67</sup> Anti-tTG antibodies binding to tissue transglutaminase in the thyroid gland have been suggested as an etiological factor.<sup>68</sup>

The prevalence of hypothyroidism is reported to be 2–13% in patients with CD in several studies equating to about a 3–4-fold higher risk (Table 2) than in the general population or controls.<sup>25,69,70</sup> Individual studies have conflicting results with regard to whether the risk of hyperthyroidism is increased.<sup>26,69,70</sup> The largest study to date (which included 14,021 CD patients) was conducted in Sweden.<sup>26</sup> This analysis found the risk of hypothyroidism, thyroiditis and hyperthyroidism to be increased (Table 2) by 2.9–4.4-fold compared to controls.<sup>26</sup> The risk of thyroid dysfunction was highest in children with CD.<sup>26</sup> Other studies have confirmed an increased risk of hypothyroidism in children and young adults with CD.<sup>71</sup> Multiple studies suggest an increased risk of positive thyroid antibodies with normal thyroid

function among CD patients as well.<sup>25,70</sup> Work in patients without CD suggests that such antibody positive patients tend to have an increased risk for developing future thyroid dysfunction.<sup>72</sup>

Longitudinal data regarding the effect of GFD upon thyroid function are limited.<sup>70,73</sup> One study concluded that strict adherence to a GFD may “normalize subclinical hypothyroidism”<sup>70</sup> while the other did not.<sup>71</sup> In the first, there was, however, no comparison to controls without CD, making it impossible to determine whether normalization of thyroid function was related to treatment or the natural history of subclinical hypothyroidism, which tends to resolve spontaneously in a sizable percentage of patients.<sup>74</sup>

A recent meta-analysis confirmed that patients with CD are at 3-fold increased risk for thyroid disease (defined as hypo- or hyperthyroidism or positive thyroid antibodies) compared to controls.<sup>25</sup> In this analysis, risk of hypothyroidism and euthyroidism with positive antibodies was increased by 3.38 (95% CI 2.88–6.56) and 4.35 (95% CI 2.88–6.56) times, respectively; in contrast, there was no increased risk of hyperthyroidism and risk of thyroid disease was not lower in those who were on a GFD compared to those who were not.<sup>25</sup>

Given the increased risk for hypothyroidism demonstrated in these studies, it is reasonable to screen patients with CD for thyroid dysfunction. Both the American Thyroid Association and the American Association of Clinical Endocrinologists recommend measurement of thyroid function in those with autoimmune diseases.<sup>75</sup> On the contrary, the prevalence of CD is also increased in those with AIT. Some experts recommend screening for CD in those with AIT, though this is not uniformly endorsed by guidelines.<sup>76,77</sup> Suspicion for CD should be raised in patients with hypothyroidism who are compliant with levothyroxine but refractory to treatment, as CD can cause malabsorption of levothyroxine.<sup>78</sup> One study suggested that hypothyroid patients requiring higher daily doses of levothyroxine are more likely to have CD. The authors of this study recommended hypothyroid patients requiring  $\geq 125$  mcg/day of levothyroxine undergo serologic testing for CD.<sup>79</sup>



### **Type 1 diabetes mellitus (T1DM)**

The same mechanisms that predispose CD patients to AIT are thought to contribute to the development of T1DM. The clinical features of new-onset T1DM are shown in Table 3. The prevalence of biopsy-diagnosed CD in children with T1DM ranges from 1.6% to 12.3%.<sup>80,81</sup> A 2014 systematic review with a meta-analysis identified 27 English-language studies, each containing over 100 patients with T1DM, who underwent screening for CD with confirmatory duodenal biopsy.<sup>82</sup> Among 26,605 type I diabetics, the estimated pooled prevalence of CD was 6% (95% CI 5.0–6.9), though there was significant heterogeneity which was not explained by subgroup analyses.<sup>82</sup> CD prevalence decreased with age at testing, with the highest CD prevalence among children (6.2%).<sup>82</sup>

Due to this elevated risk, screening for CD among patients with T1DM is recommended by some societies and guidelines;<sup>1,83</sup> however, the ideal timing for screening has not been determined. As a result of varying recommendations about CD screening in T1DM, practice patterns vary.<sup>84</sup> In a systematic review of 9 longitudinal studies containing 11,157 children with T1DM, 43.4 patients per 1,000 patient-years were diagnosed at 1 year, 32.8 at 2 years, and 20.1 at 5 years.<sup>85</sup> The authors concluded that screening for CD should be considered at T1DM diagnosis and within 2 and 5 years thereafter.<sup>85</sup> In the same study, investigators found 7% of CD diagnoses occurred before the diagnosis of T1DM.<sup>85</sup> Few studies, however, specifically evaluated the risk of T1DM in patients with CD. A Swedish prospective study, showed that children with CD are at increased risk (hazard ratio [HR] 2.4) for the subsequent development of T1DM before age 20 (Table 2), though the absolute risk was low.<sup>27</sup> A later Italian study did not confirm this but was much smaller and may have been underpowered.<sup>71</sup>

The benefit of a GFD on glycemic control and cardiovascular risk in those with coexisting T1DM and CD is unclear, as there are few longitudinal studies. One risk factor that has been shown to fairly consistently improve with institution of a GFD is high-density lipoprotein

(HDL); this may be because the intestine is a major source of apo A1.<sup>86,87</sup> Indeed, one study reported that both glycosylated hemoglobin (HbA1c) and HDL improved in GFD-adherent compared to non-adherent patients one year after GFD initiation.<sup>88</sup> It is unclear, however, if this is due to a GFD or better medical compliance in general among adherers. In contrast, another study showed an increase in HDL, but not lower HbA1c, 1 year after GFD adherence.<sup>89</sup> Studies also suggest there is an increased risk of long-term diabetes-related complications, including mortality and microvascular disease in those with T1DM and CD compared to those with T1DM alone, though GFD adherence was not assessed.<sup>90–92</sup>

### **Type 2 diabetes (T2DM)**

The literature exploring the relationship between CD and T2DM is limited, but most data indicate no increased risk for T2DM and a possible protective effect of CD. A 1998 study of 745 patients with non-insulin-dependent diabetes found CD in 0.27%.<sup>93</sup> In 2013, a US study among 840 CD patients demonstrated a lower prevalence of non-insulin-dependent diabetes (3.1% vs. 9.6%) and metabolic syndrome compared to matched controls (3.5 vs. 12.7%).<sup>94</sup> These results remained significant even when adjusting for BMI and smoking.<sup>94</sup> The authors suggest that CD patients may be protected against the development of non-insulin-dependent diabetes, though a possible mechanism remains to be determined.<sup>94</sup> In contrast, a Finnish study reported the prevalence of T2DM was 2.8% in 1,358 patients with CD, which was similar to population-based values.<sup>95</sup> However, a 2017 study evaluating the presence of IgA-tTG with confirmatory biopsy for CD in patients with T2DM on insulin therapy with poor glycemic control found CD in 1.45%, which is slightly higher than the 1% prevalence of CD worldwide.<sup>96</sup> Patients with T2DM who had lower C-peptide values also had higher levels of IgA-tTG.<sup>96</sup> The authors suggested that T2DM patients with poor glycemic control on insulin therapy who have low C-peptide levels may benefit from CD testing.<sup>96</sup>

## Adrenal disease

Autoimmune adrenal (Addison's) disease has been associated with CD in a handful of studies. The clinical features of primary adrenal insufficiency are shown in Table 3. A large Swedish study found that patients with CD were at an astonishing 11.4 times the risk for developing Addison's disease (Table 2); increased risk was observed in both children and adults.<sup>28</sup> The prevalence in CD patients was 0.25% versus 0.02% in those without CD. Most other studies have assessed the prevalence of CD in patients with Addison's disease. One study found 12.2% of patients with Addison's disease had CD.<sup>97</sup> In a Norwegian study, the prevalence of CD among 76 patients with Addison's disease identified via a registry was 7.9%.<sup>98</sup> An Italian study suggested a CD rate of 5.9% in 17 patients with Addison's disease.<sup>99</sup> Similar mechanisms predisposing to other autoimmune endocrine conditions in patients with CD likely also affect risk for autoimmune adrenal disease. In one study, those with Addison's disease and CD were all positive for the CD-associated HLA haplotype DR3-DQ2.<sup>98</sup>

## Hypogonadism and infertility

### *Female hypogonadism, reproductive disorders, and obstetric complications*

CD has been associated with female reproductive disorders, including delayed menarche, early menopause, and amenorrhea/hypogonadism.<sup>100</sup> The clinical features of female hypogonadism are shown in Table 3. Patients with CD have been reported to have later mean age of menarche compared to those with irritable bowel syndrome (IBS;  $13.7 \pm 1.5$  vs.  $12.8 \pm 1.6$  years), though age at menopause did not differ.<sup>101</sup> Secondary amenorrhea was present in 28% of women with CD, but none of those with IBS.<sup>101</sup> Subgroup analyses suggested some relationship to nutritional status. In contrast, another study found no difference in menarche age between women who had menarche before vs. after the diagnosis of CD, nor did menarche age differ compared with their mothers or healthy controls.<sup>102</sup> A 2011 study suggested that compared to healthy controls, women with untreated CD, had lower BMI and a shorter fertile

lifespan due to later menarche and earlier menopause.<sup>103</sup> In 2014, another study found that 24% of women with CD had dysfunctional uterine bleeding compared with 10% of controls; 83.3% of patients had normalization of menses after 3 months of a GFD.<sup>104</sup> Although studies are small and mostly based on historical information, they suggest CD may be associated with later menarche and amenorrhea; further research is needed in this area.

Infertility has also been reported to be a feature of CD. A 2016 meta-analysis reported a pooled CD prevalence of 2.3% (95% CI 1.4–3.5) among 884 women with infertility and a 3.2% (95% CI 2.0–4.1) prevalence among 623 women with unexplained infertility.<sup>105</sup> Compared to controls, women with infertility had 3.5 times the odds of having CD (95% CI 1.3–9.0) and women with unexplained infertility had 6 times the odds of having CD (95% CI 2.4–14.6).<sup>105</sup> In contrast, no increased risk of infertility was found in two large population-based cohort studies of women with CD, one performed among 1,521 women in the UK (HR 1.01, 95% CI 0.91–1.14) and the other performed among 11,495 women with CD in Sweden (HR 1.03, 95% CI 1.01–1.05).<sup>29,30</sup> However, as shown in Table 2, the latter study reported decreased fertility 0–2 years before the diagnosis of CD (HR 0.63; 95% CI 0.57–0.70).<sup>30</sup> Similarly, a 2014 meta-analysis reported 3.09 times higher odds of undiagnosed CD among women with infertility (95% CI 1.74–5.49), though no difference in the odds of diagnosed CD existed (OR 0.99, 95% CI 0.86–1.13).<sup>32</sup> These data suggest that women with infertility, particularly if unexplained, should be screened for CD. Because available data tend to suggest that there is no increased infertility in those with diagnosed CD, they imply that treatment of CD may restore reproductive function. Prospective longitudinal studies are needed, however, to confirm this.

In women with CD who become pregnant, spontaneous abortion and several obstetrical complications have been reported to be higher in comparison to those without CD.<sup>106,107</sup> A recent study indicated those with recurrent pregnancy loss, were more likely have CD-associated haplotype DQ2/DQ8 compared to controls.<sup>108</sup> Obstetrical complications have been assessed

recently in a 2016 meta-analysis that included 10 cohort studies and 4.8 million women.<sup>107</sup> It indicated women with CD had a significantly higher risk for the development of preterm birth, intrauterine growth restriction, still birth, low birthweight and small for gestational age infants.<sup>107</sup> Risk for preeclampsia was not increased. Subgroup analyses suggested that women with diagnosed and treated CD had a lower risk for preterm birth compared to those with undiagnosed and untreated CD, indicating that adherence to a GFD could alter the risk of this complication.

The mechanisms causing reproductive dysfunction in CD have been inadequately investigated to date and remain to be elucidated. Most studies have not assessed hormonal regulators of reproductive function. Nutritional deficits may play a role. Zinc and selenium deficiencies have been linked to impaired synthesis and secretion of FSH and LH,<sup>109</sup> and have been reported in children with untreated CD.<sup>110,111</sup> Degree of malnutrition and low weight may contribute to delayed menarche and secondary amenorrhea in CD, as they do in other conditions.<sup>101</sup> Only one study to date has evaluated pituitary gland function in women with CD. This 2018 study failed to show altered values of LH, FSH, and prolactin in 46 women with CD, but did report decreased anti-Mullerian hormone (AMH), an indicator of declining ovarian function and follicular reserve.<sup>112</sup> Given this study's small sample size,<sup>112</sup> larger studies evaluating gonadotropins and sex hormones in women with CD are warranted. Other studies suggest that infertility may be caused by anti-tTG antibodies binding to trophoblastic cells and/or endometrial endothelial cells leading to placental damage.<sup>113–115</sup> These mechanisms could explain the increased risk of miscarriage, low birth weight, and preterm delivery demonstrated in pregnant women with CD.<sup>115</sup>

### **Male hypogonadism**

Fewer studies have focused on male reproductive disorders in CD, but case reports of hypogonadism can be found as early as the 1950s.<sup>116</sup> A small study from the 1980s comparing 28 men with CD to 19 men with Crohn's disease reported increased rates of hypogonadism (7 vs. 0%) and

sexual dysfunction in CD compared to controls.<sup>117</sup> The clinical features of male hypogonadism are shown in Table 3. Males with CD also had abnormal sperm morphology and motility. Sperm morphology improved after gluten withdrawal.<sup>117</sup> Additionally, infertility was reported in 19% of males with CD in this study.<sup>117</sup> Sperm impairment was unrelated to nutritional deficiencies, including B12 and folate, or to infertility.

As shown in Table 2, results from a large population-based cohort study of 7,121 Swedish men with biopsy-verified CD from 2011, however, are reassuring, as there was no significant increase in infertility (HR of 1.02 (95% CI 0.99–1.04)).<sup>31</sup> Similar results were reported in a recent meta-analysis (Table 2).<sup>32</sup> Rates of CD among men reporting infertility have also been investigated. Two studies of couples with infertility found a CD prevalence of 0.39–1% among males,<sup>118,119</sup> while a third found a higher prevalence of 6%.<sup>120</sup> A meta-analysis of two of these studies,<sup>118,120</sup> however, reported no increased odds of CD in this group (OR 1.63, 95% CI 0.60–4.46). The majority of these patients were diagnosed by serology only.<sup>32</sup>

Small studies of men with CD from the 1970s and 1980s have reported androgen resistance, characterized by increased basal FSH and LH,<sup>121–123</sup> increased plasma testosterone and free testosterone index,<sup>122,123</sup> elevated plasma SHBG,<sup>121</sup> exaggerated gonadotropin responses to LH-releasing hormone,<sup>121</sup> and reduced dihydrotestosterone (DHT) in men with CD.<sup>122,123</sup> These abnormalities were not present in controls with Crohn's disease who had similar nutritional status.<sup>122,123</sup> Available data suggest that treatment with a GFD alleviates androgen insensitivity; in several studies, laboratory abnormalities (with the exception of increased basal and stimulated FSH values) returned to normal with healing of jejunal morphology after initiation of a GFD.<sup>121–123</sup> Elevated prolactin values have also been reported in men<sup>121</sup> and children with untreated CD;<sup>124,125</sup> however, these values were only moderately elevated and have not been associated with impotence or infertility.<sup>121</sup> Despite no apparent risk of male infertility in CD overall, further studies are needed to verify if there is any association between CD and pituitary or peripheral regulation of gonadal function.

### **Growth hormone deficiency and growth**

Short stature or failure to thrive can be the presenting sign of CD in children.<sup>126–128</sup> Most children with CD and short stature display “catch up” growth when a GFD is instituted.<sup>128</sup> A subset of children may also have transient functional GHD, growth hormone resistance or low insulin-like growth factor-1 (IGF-1) that reverses with institution of a GFD.<sup>129</sup> The clinical features of GHD are shown in Table 3. One study indicated that among children with short stature, 0.63% had CD and 0.23% had both CD and GHD.<sup>130</sup> Lack of an increase in growth velocity after adherence to a GFD should raise suspicion that GHD may be congenital and/or permanent.<sup>128,131,132</sup> One study indicated that those who responded to a GFD with an appropriate increase in growth velocity had more delayed bone age than nonresponders, suggesting this could be helpful in determining who may be most likely to respond.<sup>132</sup> Treatment with growth hormone in combination with a GFD can increase growth rate in children with GHD and CD.<sup>128,130,133</sup> Final adult height in CD patients is typically similar to that of the general population, unless there is a delay in diagnosis.<sup>134,135</sup>

One study suggested that 25% of adults with CD may have complete or partially impaired growth hormone secretion that is independent from diagnosis and implementation of a GFD.<sup>136</sup> All those with GHD were men and none had anti-pituitary antibodies (APA).<sup>136</sup> Conversely, a high percentage (42%) of APA antibodies has been reported in children newly diagnosed with CD compared to only 2% of controls without CD.<sup>137</sup> IGF-1 was lower in patients who were APA positive compared to those who were APA negative and height was positively correlated with IGF-1.<sup>137</sup>

### **Autoimmune polyendocrine syndromes**

CD may occur as part of an autoimmune polyendocrine syndrome (APS), a rare collection of disorders characterized by the autoimmune failure of two or more endocrine glands. APS type 1 (APS-1), also known as autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy (APECED), is a rare autosomal recessive condition due to a mutation in the autoimmune regulator (*AIRE*)

gene; it is characterized by the development of two of three manifestations, including chronic mucocutaneous candidiasis, hypoparathyroidism, and Addison’s disease, during childhood.<sup>138</sup> CD is not a feature of APS-1.

On the other hand, APS type 2 (APS-2), a more common, polygenic condition presenting in adolescence or adulthood, is characterized (Table 3) by at least two of three endocrinopathies: Addison’s disease, T1DM, and thyroid disease.<sup>138,139</sup> CD can be a feature of APS-2 along with other non-endocrine autoimmune conditions such as alopecia, vitiligo, and pernicious anemia.<sup>138–142</sup> Primary ovarian failure has also been described. APS-2 is a clinical diagnosis supported by the presence of auto-antibodies against thyroid peroxidase, glutamic acid decarboxylase 65, and to 21-hydroxylase.<sup>138</sup> Though some suggest further subcategorizing APS-2 without adrenal involvement as APS-3 and CD has been reported in the literature in association with “APS-3”, it is not clearly recognized as a separate entity.<sup>143,144</sup>

Patients with concomitant APS-2 and CD have been found to harbor variants in haplotypes DR3-DQ2 and DR4-DQ8 which have also been associated with T1DM, AIT, and Addison’s disease in other studies, indicating a common genetic predisposition.<sup>138,145–147</sup> A 2016 study found that patients with “APS-3” had a higher frequency of positive tTG antibodies (36%) compared to patients with T1DM or AIT alone, though this study only tested for CD autoimmunity and not biopsy-confirmed CD.<sup>148</sup>

As noted in prior sections, CD can occur in conjunction with autoimmune failure of one or more endocrine glands (including the thyroid, pancreas or adrenal), outside of the context of an recognized APS syndrome. Further, the coexistence of CD in a patient with one autoimmune endocrinopathy may increase the risk for involvement of multiple endocrine glands. For example, a recent study suggests the risk of AIT was increased in patients with T1DM and CD (HR 1.67, 95% CI 1.32–2.11) compared to those with T1DM alone. The highest risk occurred in those with CD for 10 years or more (HR 2.22, 95% CI 1.49–3.23).<sup>149</sup>

In addition to the associations of CD with AIT, T1DM, and Addison’s disease, CD has rarely been described in association with idiopathic hypoparathyroidism and autoimmune hypophysitis in

separate isolated reports or case series.<sup>150–154</sup> One study suggested a potential mechanism linking hypoparathyroidism and untreated CD cross-reactivity between CD-specific EMA antibodies and parathyroid tissue leading to parathyroid atrophy.<sup>155</sup> This mechanism, however, has not been confirmed. Further, a more recent systematic investigation reported no increased risk of CD in patients with idiopathic hypoparathyroidism.<sup>150</sup> The association between CD and autoimmune hypophysitis has not been well defined to date.

### **Summary and key points**

Patients with CD are at increased risk for metabolic bone disease, fractures, and a number of less frequent autoimmune and non-autoimmune endocrine disorders. The patient described in the Case Report illustrates several endocrine conditions associated with CD and underscores key points evident from review of the literature on the endocrine complications of CD. Importantly, the diagnosis of CD is often delayed in those presenting with extra-intestinal manifestations rather than classic symptoms of malabsorption.<sup>1,2,7</sup> Early diagnosis may be facilitated by adherence to published guidelines that include fatigue and weight loss alone as indications to test for CD.<sup>1</sup>

Osteoporosis is thought to be a common finding in CD and its evaluation may lead to diagnosis of CD, as described in the Case Report. Consistent with the literature on metabolic bone disease in CD,<sup>14,19,37</sup> the pathogenesis of bone loss in this case involved calcium (if not vitamin D) malabsorption leading to osteomalacia (presumably), compensatory secondary hyperparathyroidism, and increased bone turnover. Although all laboratory abnormalities related to the patient's metabolic bone disease resolved with duodenal mucosal healing on GFD, restoration of BMD was incomplete, as shown in numerous studies (though antiepileptic therapy may have contributed to residual reduced BMD in this case).<sup>156</sup> As is often the case in clinical practice, pharmacological osteoporosis treatment was ultimately needed for the patient described.

The patient's presentation and course were also marked by clinical and laboratory findings consistent with the less frequent CD complication of incomplete androgen insensitivity, with partial resolution on GFD

as reported in several studies of men with CD;<sup>119–121</sup> as with persistent osteoporosis in this case, the mild, residual elevation of SHBG on GFD may have been attributed to treatment with phenytoin, which is known to increase circulating SHBG levels.<sup>157</sup> Recent studies also suggest other reproductive consequences of CD, not seen in this case, including hypogonadism, female infertility and preterm labor in women. The case highlights that testing for CD in those with osteoporosis or reproductive dysfunction should be considered due to the possibility of improvement after the diagnosis of CD and institution of a GFD. Conclusive evidence regarding the resolution or improvement of the endocrine manifestations of CD with institution of a GFD, however, is lacking and studies are needed that objectively address this issue.

Finally, the patient had evidence of CD-associated endocrine gland autoimmunity, in this case AIT<sup>68,69</sup> and thyroid parenchymal heterogeneity but no overt thyroid dysfunction. Strong epidemiological associations between CD and several autoimmune endocrinopathies, including not only AIT but also T1DM and Addison's disease, have been established by recent large prospective cohort studies. Common genetic variants and HLA haplotypes are thought to predispose to both CD and endocrine autoimmunity, though much of the available data is associative in nature and exact mechanisms remain unclear.

In conclusion, the endocrine complications of CD – many manifested by the patient reported herein – are not uncommon but remain incompletely understood. Given most of the data pertaining to the endocrine manifestations of CD have been collected in Caucasian populations, it is unclear whether risk estimates are applicable to non-Caucasians. Studies assessing ethnic and racial differences in risk for endocrine disease are needed. Research into underlying mechanisms and associations is also warranted and will likely inform improvement in clinical diagnosis and treatment. Unfortunately, the diagnosis of CD in those manifesting only endocrine dysfunction may be delayed. Screening for CD may be appropriate in certain high prevalence groups such as those with osteoporosis, hypothyroidism, T1DM or multiple endocrinopathies and is recommended by some experts.<sup>158</sup> Likewise, gastroenterologists who care for CD patients must have a high degree of suspicion for

possible concomitant endocrine disorders, particularly those that are frequent, such as osteoporosis, vitamin D deficiency, and hypothyroidism.

## Key points

- (1) Patients with CD are at increased risk for metabolic bone disease and a number of less frequent autoimmune and non-autoimmune endocrine disorders.
- (2) Diagnosis of CD is often delayed in those presenting with extra-intestinal manifestations.
- (3) Screening for CD may be appropriate in certain high prevalence groups such as patients with osteoporosis, hypothyroidism, T1DM, and multiple endocrinopathies and is recommended by some experts.
- (4) The link between autoimmune endocrinopathies and CD is incompletely understood but is likely due in part to HLA genotypes predisposing to both conditions.

## Acknowledgments

The authors express their appreciation to the patient and his wife for their gracious consent and collaborative assistance in presentation of his medical history as a Case Report.

## Declaration of Interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

## References

1. Lebwohl B, Sanders DS, Green PHR. Coeliac disease. *Lancet*. 2018;391(10115):70–81. doi:10.1016/S0140-6736(17)31796-8.
2. Leffler DA, Green PH, Fasano A. Extraintestinal manifestations of coeliac disease. *Nat rev Gastroenterol hepatol*. 2015;12(10):561–571. doi:10.1038/nrgastro.2015.131.
3. Mustalahti K, Catassi C, Reunanen A, et al. The prevalence of celiac disease in Europe: results of a centralized, international mass screening project. *Ann Med*. 2010;42(8):587–595. doi:10.3109/07853890.2010.505931.
4. Singh P, Arora A, Strand TA, et al. Global prevalence of celiac disease: systematic review and meta-analysis. *Clin gastroenterol hepatol*. 2018;16(6):823–36 e2. doi:10.1016/j.cgh.2017.06.037.
5. Choung RS, Ditah IC, Nadeau AM, et al. Trends and racial/ethnic disparities in gluten-sensitive problems in the United States: findings from the National Health and Nutrition Examination Surveys from 1988 to 2012. *Am J Gastroenterol*. 2015;110(3):455–461. doi:10.1038/ajg.2015.8.
6. Makharia GK, Verma AK, Amarchand R, et al. Prevalence of celiac disease in the northern part of India: a community based study. *J Gastroenterol Hepatol*. 2011;26(5):894–900. doi:10.1111/j.1440-1746.2010.06606.x.
7. Lohi S, Mustalahti K, Kaukinen K, et al. Increasing prevalence of coeliac disease over time. *Aliment Pharmacol Ther*. 2007;26(9):1217–1225. doi:10.1111/j.1365-2036.2007.03502.x.
8. Murray JA, Van Dyke C, Plevak MF, Dierkhising RA, Zinsmeister AR, Melton LJ 3rd. Trends in the identification and clinical features of celiac disease in a North American community, 1950–2001. *Clin gastroenterol hepatol*. 2003;1(1):19–27. doi:10.1053/jcgh.2003.50004.
9. Choung RS, Unalp-Arida A, Ruhl CE, Brantner TL, Everhart JE, Murray JA. Less hidden celiac disease but increased gluten avoidance without a diagnosis in the United States: findings from the national health and nutrition examination surveys from 2009 to 2014. *Mayo Clin Proc*. 2016;92(1): 30–38. doi:10.1016/j.mayocp.2016.10.012.
10. Dixit R, Lebwohl B, Ludvigsson JF, Lewis SK, Rizkalla-Reilly N, Green PH. Celiac disease is diagnosed less frequently in young adult males. *Dig Dis Sci*. 2014;59(7):1509–1512. doi:10.1007/s10620-014-3025-6.
11. Kagnoff MF. Celiac disease: pathogenesis of a model immunogenetic disease. *J Clin Invest*. 2007;117(1):41–49. doi:10.1172/JCI30253.
12. Olivares M, Neef A, Castillejo G, et al. The HLA-DQ2 genotype selects for early intestinal microbiota composition in infants at high risk of developing coeliac disease. *Gut*. 2015;64(3):406–417. doi:10.1136/gutjnl-2014-306931.
13. Hopper AD, Hadjivassiliou M, Hurlstone DP, et al. What is the role of serologic testing in celiac disease? A prospective, biopsy-confirmed study with economic analysis. *Clin Gastroenterol Hepatol*. 2008;6(3):314–320. doi:10.1016/j.cgh.2007.12.008.
14. Husby S, Koletzko S, Korponay-Szabó IR, et al. European society for pediatric gastroenterology, hepatology, and nutrition guidelines for the diagnosis of coeliac disease. *J Pediatr Gastroenterol Nutr*. 2012;54(1):136–160. doi:10.1097/MPG.0b013e31821a23d0.
15. Zanchetta MB, Longobardi V, Bai JC. Bone and celiac disease. *Curr Osteoporos Rep*. 2016;14(2):43–48. doi:10.1007/s11914-016-0304-5.
16. Larussa T, Suraci E, Nazionale I, Abenavoli L, Imeneo M, Luzza F. Bone mineralization in celiac disease. *Gastroenterol Res Pract*. 2012;2012:198025. doi:10.1155/2012/585674.
17. Meyer D, Stavropoulos S, Diamond B, Shane E, Green PH. Osteoporosis in a north american adult population

- with celiac disease. *Am J Gastroenterol.* 2001;96(1):112–119. doi:10.1111/j.1572-0241.2001.03507.x.
18. Walters JR, Banks LM, Butcher GP, Fowler CR. Detection of low bone mineral density by dual energy x ray absorptiometry in unsuspected suboptimally treated coeliac disease. *Gut.* 1995;37:220–224.
  19. Motta ME, Faria ME, Silva GA. Prevalence of low bone mineral density in children and adolescents with celiac disease under treatment. *Sao Paulo Medical Journal = Revista Paulista De Medicina.* 2009;127:278–282.
  20. Laszkowska M, Mahadev S, Sundstrom J, et al. Systematic review with meta-analysis: the prevalence of coeliac disease in patients with osteoporosis. *Aliment Pharmacol Ther.* 2018. doi:10.1111/apt.14911.
  21. Corazza GR, Di Stefano M, Maurino E, Bai JC. Bones in coeliac disease: diagnosis and treatment. *Best Practice Res Clin Gastroenterol.* 2005;19(3):453–465. doi:10.1016/j.bpg.2005.01.002.
  22. Kamycheva E, Goto T, Camargo CA Jr. Celiac disease is associated with reduced bone mineral density and increased FRAX scores in the US national health and nutrition examination survey. *Osteoporos Int.* 2017;28(3):781–790. doi:10.1007/s00198-016-3791-4.
  23. Ludvigsson JF, Michaelsson K, Ekbom A, Montgomery SM. Coeliac disease and the risk of fractures - a general population-based cohort study. *Aliment Pharmacol Ther.* 2007;25(3):273–285. doi:10.1111/j.1365-2036.2006.03203.x.
  24. Heikkila K, Pearce J, Maki M, Kaukinen K. Celiac disease and bone fractures: a systematic review and meta-analysis. *J Clin Endocrinol Metab.* 2015;100(1):25–34. doi:10.1210/jc.2014-1858.
  25. Sun X, Lu L, Yang R, Li Y, Shan L, Wang Y. Increased incidence of thyroid disease in patients with celiac disease: a systematic review and meta-analysis. *PLoS One.* 2016;11(12):e0168708. doi:10.1371/journal.pone.0168708.
  26. Elfstrom P, Montgomery SM, Kampe O, Ekbom A, Ludvigsson JF. Risk of thyroid disease in individuals with celiac disease. *J Clin Endocrinol Metab.* 2008;93(10):3915–3921. doi:10.1210/jc.2008-0798.
  27. Ludvigsson JF, Ludvigsson J, Ekbom A, Montgomery SM. Celiac disease and risk of subsequent type 1 diabetes: a general population cohort study of children and adolescents. *Diabetes Care.* 2006;29(11):2483–2488. doi:10.2337/dc06-0794.
  28. Elfstrom P, Montgomery SM, Kampe O, Ekbom A, Ludvigsson JF. Risk of primary adrenal insufficiency in patients with celiac disease. *J Clin Endocrinol Metab.* 2007;92(9):3595–3598. doi:10.1210/jc.2007-0960.
  29. Tata LJ, Card TR, Logan RF, Hubbard RB, Smith CJ, West J. Fertility and pregnancy-related events in women with celiac disease: a population-based cohort study. *Gastroenterology.* 2005;128:849–855.
  30. Zugna D, Richiardi L, Akre O, Stephansson O, Ludvigsson JF. A nationwide population-based study to determine whether coeliac disease is associated with infertility. *Gut.* 2010;59(11):1471–1475. doi:10.1136/gut.2010.219030.
  31. Zugna D, Richiardi L, Akre O, Stephansson O, Ludvigsson JF. Celiac disease is not a risk factor for infertility in men. *Fertility and Sterility.* 2011;95(5):1709–13e1-3. doi:10.1016/j.fertnstert.2011.01.132.
  32. Lasa JS, Zubiaurre I, Soifer LO. Risk of infertility in patients with celiac disease: a meta-analysis of observational studies. *Arquivos de gastroenterologia.* 2014;51:144–150.
  33. Moreno ML, Vazquez H, Mazure R, et al. Stratification of bone fracture risk in patients with celiac disease. *Clin gastroenterol hepatol.* 2004;2(2):127–134.
  34. Lebwohl B, Michaelsson K, Green PH, Ludvigsson JF. Persistent mucosal damage and risk of fracture in celiac disease. *J Clin Endocrinol Metab.* 2014;99(2):609–616. doi:10.1210/jc.2013-3164.
  35. Vestergaard P, Mosekilde L. Fracture risk in patients with celiac Disease, Crohn's disease, and ulcerative colitis: a nationwide follow-up study of 16,416 patients in Denmark. *Am J Epidemiol.* 2002;156:1–10.
  36. Davie MW, Gaywood I, George E, et al. Excess non-spine fractures in women over 50 years with celiac disease: a cross-sectional, questionnaire-based study. *Osteoporos Int.* 2005;16(9):1150–1155. doi:10.1007/s00198-004-1822-z.
  37. Stein EM, Rogers H, Leib A, et al. Abnormal skeletal strength and microarchitecture in women with celiac disease. *J Clin Endocrinol Metab.* 2015;100(6):2347–2353. doi:10.1210/jc.2015-1392.
  38. Zanchetta MB, Costa F, Longobardi V, et al. Significant bone microarchitecture impairment in premenopausal women with active celiac disease. *Bone.* 2015;76:149–157. doi:10.1016/j.bone.2015.03.005.
  39. Bianchi ML, Bardella MT. Bone in celiac disease. *Osteoporos Int.* 2008;19(12):1705–1716. doi:10.1007/s00198-008-0624-0.
  40. Selby PL, Davies M, Adams JE, Mawer EB. Bone loss in celiac disease is related to secondary hyperparathyroidism. *J Bone Miner Res.* 1999;14(4):652–657. doi:10.1359/jbmr.1999.14.4.652.
  41. Keaveny AP, Freaney R, McKenna MJ, Masterson J, O'Donoghue DP. Bone remodeling indices and secondary hyperparathyroidism in celiac disease. *Am J Gastroenterol.* 1996;91:1226–1231.
  42. Ludvigsson JF, Kampe O, Lebwohl B, Green PH, Silverberg SJ, Ekbom A. Primary hyperparathyroidism and celiac disease: a population-based cohort study. *J Clin Endocrinol Metab.* 2012;97(3):897–904. doi:10.1210/jc.2011-2639.
  43. Zylberberg HM, Lebwohl B, RoyChoudhury A, Walker MD, Green PHR. Predictors of improvement in bone mineral density after celiac disease diagnosis. *Endocrine.* 2018;59(2):311–318. doi:10.1007/s12020-017-1488-x.
  44. Tilg H, Moschen AR, Kaser A, Pines A, Dotan I. Gut, inflammation and osteoporosis: basic and clinical concepts. *Gut.* 2008;57(5):684–694. doi:10.1136/gut.2006.117382.

45. Fiore CE, Pennisi P, Ferro G, et al. Altered osteoprotegerin/RANKL ratio and low bone mineral density in celiac patients on long-term treatment with gluten-free diet. *Hormone and Metabolic Research = Hormon- Und Stoffwechselforschung = Hormones Et Metabolisme*. 2006;38(6):417–422. doi:10.1055/s-2006-944548.
46. Riches PL, McRorie E, Fraser WD, Determann C. Osteoporosis associated with neutralizing autoantibodies against osteoprotegerin. *N Engl J Med*. 2009;361:1459–1465.
47. Pantaleoni S, Luchino M, Adriani A, et al. Bone mineral density at diagnosis of celiac disease and after 1 year of gluten-free diet. *Sci World J*. 2014;2014:173082. doi:10.1155/2014/173082.
48. Grace-Farfaglia P. Bones of contention: bone mineral density recovery in celiac disease—a systematic review. *Nutrients*. 2015;7(5):3347–3369. doi:10.3390/nu7053347.
49. Valdimarsson T, Lofman O, Toss G, Strom M. Reversal of osteopenia with diet in adult coeliac disease. *Gut*. 1996;38:322–327.
50. Szymczak J, Bohdanowicz-Pawlak A, Waszczuk E, Jakubowska J. Low bone mineral density in adult patients with coeliac disease. *Endokrynologia Polska*. 2012;63:270–276.
51. Kotze LM, Skare T, Vinholi A, Jurkonis L, Nisihara R. Impact of a gluten-free diet on bone mineral density in celiac patients. *Revista espanola de enfermedades digestivas: organo oficial de la Sociedad Espanola de Patologia Digestiva*. 2016;108(2):84–88. doi:10.17235/reed.2015.3953/2015.
52. Zanchetta MB, Longobardi V, Costa F, et al. Impaired bone microarchitecture improves after one year on gluten-free diet: a prospective longitudinal HRpQCT study in women with celiac disease. *J Bone Miner Res*. 2017;32(1):135–142. doi:10.1002/jbmr.2922.
53. Ciacci C, Maurelli L, Klain M, et al. Effects of dietary treatment on bone mineral density in adults with celiac disease: factors predicting response. *Am J Gastroenterol*. 1997;92(6):992–996.
54. Bai JC, Gonzalez D, Mautalen C, et al. Long-term effect of gluten restriction on bone mineral density of patients with coeliac disease. *Aliment Pharmacol Ther*. 1997;11(1):157–164.
55. Casella S, Zanini B, Lanzarotto F, Villanacci V, Ricci C, Lanzini A. Celiac disease in elderly adults: clinical, serological, and histological characteristics and the effect of a gluten-free diet. *J Am Geriatr Soc*. 2012;60(6):1064–1069. doi:10.1111/j.1532-5415.2012.03997.x.
56. Kempainen T, Kroger H, Janatuinen E, et al. Bone recovery after a gluten-free diet: a 5-year follow-up study. *Bone*. 1999;25(3):355–360.
57. Sategna-Guidetti C, Grosso SB, Grosso S, et al. The effects of 1-year gluten withdrawal on bone mass, bone metabolism and nutritional status in newly-diagnosed adult coeliac disease patients. *Aliment Pharmacol Ther*. 2000;14(1):35–43.
58. Larussa T, Suraci E, Imeneo M. Normal bone mineral density associates with duodenal mucosa healing in adult patients with celiac disease on a gluten-free diet. *Nutrients*. 2017;9(2):E98. doi:10.3390/nu9020098.
59. Posthumus L, Al-Toma A. Duodenal histopathology and laboratory deficiencies related to bone metabolism in coeliac disease. *Eur J Gastroenterol Hepatol*. 2017;29(8):897–903. doi:10.1097/MEG.0000000000000880.
60. Kumar M, Rastogi A, Bhadada SK, Bhansali A, Vaiphei K, Kochhar R. Effect of zoledronic acid on bone mineral density in patients of celiac disease: a prospective, randomized, pilot study. *Indian J Med Res*. 2013;138:882–887.
61. Meek SE, Nix K. Hypocalcemia after alendronate therapy in a patient with celiac disease. *Endocr Pract*. 2007;13(4):403–407. doi:10.4158/EP.13.4.403.
62. Larizza D, Calcaterra V, Klersy C, et al. Common immunogenetic profile in children with multiple autoimmune diseases: the signature of HLA-DQ pleiotropic genes. *Autoimmunity*. 2012;45(6):470–475. doi:10.3109/08916934.2012.697594.
63. Zhernakova A, Eerligh P, Barrera P, et al. CTLA4 is differently associated with autoimmune diseases in the Dutch population. *Hum Genet*. 2006;119(1–2):225. doi:10.1007/s00439-005-0087-8.
64. Tolone C, Cirillo G, Papparella A, et al. A common CTLA4 polymorphism confers susceptibility to autoimmune thyroid disease in celiac children. *Digestive Liver Disease*. 2009;41(6):385–389. doi:10.1016/j.dld.2008.09.001.
65. Djilali-Saiah I, Schmitz J, Harfouch-Hammoud E, Mougenot JF, Bach JF, Caillat-Zucman S. CTLA-4 gene polymorphism is associated with predisposition to coeliac disease. *Gut*. 1998;43:187–189.
66. Cosnes J, Cellier C, Viola S, et al. Incidence of autoimmune diseases in celiac disease: protective effect of the gluten-free diet. *Clin gastroenterol hepatol*. 2008;6(7):753–758. doi:10.1016/j.cgh.2007.12.022.
67. Toscano V, Conti FG, Anastasi E, et al. Importance of gluten in the induction of endocrine autoantibodies and organ dysfunction in adolescent celiac patients. *Am J Gastroenterol*. 2000;95(7):1742–1748. doi:10.1111/j.1572-0241.2000.02187.x.
68. Naiyer AJ, Shah J, Hernandez L, et al. Tissue transglutaminase antibodies in individuals with celiac disease bind to thyroid follicles and extracellular matrix and may contribute to thyroid dysfunction. *Thyroid*. 2008;18(11):1171–1178. doi:10.1089/thy.2008.0110.
69. Counsell CE, Taha A, Ruddell WS. Coeliac disease and autoimmune thyroid disease. *Gut*. 1994;35:844–846.
70. Sategna-Guidetti C, Volta U, Ciacci C, et al. Prevalence of thyroid disorders in untreated adult celiac disease patients and effect of gluten withdrawal: an Italian multicenter study. *Am J Gastroenterol*. 2001;96(3):751–757. doi:10.1111/j.1572-0241.2001.03617.x.
71. Canova C, Pitter G, Ludvigsson JF, et al. Celiac disease and risk of autoimmune disorders: a population-based matched birth cohort study. *J Pediatr*. 2016;174:146–152 e1. doi:10.1016/j.jpeds.2016.02.058.



72. Walsh JP, Bremner AP, Feddema P, Leedman PJ, Brown SJ, O'Leary P. Thyrotropin and thyroid antibodies as predictors of hypothyroidism: a 13-year, longitudinal study of a community-based cohort using current immunoassay techniques. *J Clin Endocrinol Metab.* 2010;95(3):1095–1104. doi:10.1210/jc.2009-1977.
73. Metso S, Hyytia-Ilmonen H, Kaukinen K, et al. Gluten-free diet and autoimmune thyroiditis in patients with celiac disease. A Prospective Controlled Study. *Scand J Gastroenterol.* 2012;47(1):43–48. doi:10.3109/00365521.2011.639084.
74. Diez JJ, Iglesias P, Burman KD. Spontaneous normalization of thyrotropin concentrations in patients with subclinical hypothyroidism. *J Clin Endocrinol Metab.* 2005;90(7):4124–4127. doi:10.1210/jc.2005-0375.
75. Garber JR, Cobin RH, Gharib H, et al. Clinical practice guidelines for hypothyroidism in adults: cosponsored by the American Association of Clinical Endocrinologists and the American Thyroid Association. *Endocr Pract.* 2012;18(6):988–1028. doi:10.4158/EP12280.GL.
76. Reilly NR, Verma R. Time to screen children with celiac disease for thyroid disease? *J Pediatr.* 2016;174:7–9. doi:10.1016/j.jpeds.2016.04.019.
77. Rubio-Tapia A, Hill ID, Kelly CP, Calderwood AH, Murray JA. American College of G. ACG clinical guidelines: diagnosis and management of celiac disease. *Am J Gastroenterol.* 2013;108(5):656–676. quiz 77. doi:10.1038/ajg.2013.79.
78. Centanni M, Benvenga S, Sachmechi I. Diagnosis and management of treatment-refractory hypothyroidism: an expert consensus report. *J Endocrinol Invest.* 2017;40(12):1289–1301. doi:10.1007/s40618-017-0706-y.
79. Zubarik R, Ganguly E, Nathan M, Vecchio J. Celiac disease detection in hypothyroid patients requiring elevated thyroid supplementation: A prospective cohort study. *Eur J Intern Med.* 2015;26(10):825–829. doi:10.1016/j.ejim.2015.09.011.
80. Hansen D, Brock-Jacobsen B, Lund E, et al. Clinical benefit of a gluten-free diet in type 1 diabetic children with screening-detected celiac disease: a population-based screening study with 2 years' follow-up. *Diabetes Care.* 2006;29(11):2452–2456. doi:10.2337/dc06-0990.
81. Poulain C, Johanet C, Delcroix C, Levy-Marchal C, Tubiana-Rufi N. Prevalence and clinical features of celiac disease in 950 children with type 1 diabetes in France. *Diabetes & Metabolism.* 2007;33(6):453–458. doi:10.1016/j.diabet.2007.06.004.
82. Elfstrom P, Sundstrom J, Ludvigsson JF. Systematic review with meta-analysis: associations between coeliac disease and type 1 diabetes. *Aliment Pharmacol Ther.* 2014;40(10):1123–1132. doi:10.1111/apt.12973.
83. Downey L, Houten R, Murch S, Longson D, Guideline Development G. Recognition, assessment, and management of coeliac disease: summary of updated NICE guidance. *BMJ.* 2015;351:h4513.
84. Simpson SM, Ciaccio EJ, Case S, et al. Celiac disease in patients with type 1 diabetes: screening and diagnostic practices. *Diabetes Educ.* 2013;39(4):532–540. doi:10.1177/0145721713487998.
85. Pham-Short A, Donaghue KC, Ambler G, Phelan H, Twigg S, Craig ME. Screening for celiac disease in type 1 diabetes: a systematic review. *Pediatrics.* 2015;136(1):e170–6. doi:10.1542/peds.2014-2883.
86. Brar P, Kwon GY, Holleran S, et al. Change in lipid profile in celiac disease: beneficial effect of gluten-free diet. *Am J Med.* 2006;119(9):786–790. doi:10.1016/j.amjmed.2005.12.025.
87. Glickman RM, Green PH. The intestine as a source of apolipoprotein A1. *Proc Natl Acad Sci U S A.* 1977;74:2569–2573.
88. Leeds JS, Hopper AD, Hadjivassiliou M, Tesfaye S, Sanders DS. High prevalence of microvascular complications in adults with type 1 diabetes and newly diagnosed celiac disease. *Diabetes Care.* 2011;34(10):2158–2163. doi:10.2337/dc11-0149.
89. Warncke K, Liptay S, Frohlich-Reiterer E, et al. Vascular risk factors in children, adolescents, and young adults with type 1 diabetes complicated by celiac disease: results from the DPV initiative. *Pediatric Diabetes.* 2016;17(3):191–198. doi:10.1111/pedi.12261.
90. Mollazadegan K, Kugelberg M, Montgomery SM, Sanders DS, Ludvigsson J, Ludvigsson JF. A population-based study of the risk of diabetic retinopathy in patients with type 1 diabetes and celiac disease. *Diabetes Care.* 2013;36(2):316–321. doi:10.2337/dc12-0766.
91. Mollazadegan K, Sanders DS, Ludvigsson J, Ludvigsson JF. Long-term coeliac disease influences risk of death in patients with type 1 diabetes. *J Intern Med.* 2013;274(3):273–280. doi:10.1111/joim.12092.
92. Rohrer TR, Wolf J, Liptay S, et al. Microvascular complications in childhood-onset type 1 diabetes and celiac disease: a multicenter longitudinal analysis of 56,514 patients from the german-Austrian DPV database. *Diabetes Care.* 2015;38(5):801–807. doi:10.2337/dc14-0683.
93. Sjoberg K, Eriksson KF, Bredberg A, Wassmuth R, Eriksson S. Screening for coeliac disease in adult insulin-dependent diabetes mellitus. *J Intern Med.* 1998;243:133–140.
94. Kabbani TA, Kelly CP, Betensky RA, et al. Patients with celiac disease have a lower prevalence of non-insulin-dependent diabetes mellitus and metabolic syndrome. *Gastroenterology.* 2013;144(5):912–7 e1. doi:10.1053/j.gastro.2013.01.033.
95. Kylokas A, Kaukinen K, Huhtala H, Collin P, Maki M, Kurppa K. Type 1 and type 2 diabetes in celiac disease: prevalence and effect on clinical and histological presentation. *BMC Gastroenterol.* 2016;16(1):76. doi:10.1186/s12876-016-0488-2.
96. Kizilgul M, Ozcelik O, Beysel S, et al. Screening for celiac disease in poorly controlled type 2 diabetes

- mellitus: worth it or not? *BMC Endocr Disord.* 2017;17(1):62. doi:10.1186/s12902-017-0212-4.
97. O'Leary C, Walsh CH, Wieneke P, et al. Coeliac disease and autoimmune Addison's disease: a clinical pitfall. *QJM.* 2002;95(2):79–82.
  98. Myhre AG, Aarsetoy H, Undlien DE, Hovdenak N, Aksnes L, Husebye ES. High frequency of coeliac disease among patients with autoimmune adrenocortical failure. *Scand J Gastroenterol.* 2003;38:511–515.
  99. Biagi F, Campanella J, Soriani A, Vailati A, Corazza GR. Prevalence of coeliac disease in Italian patients affected by Addison's disease. *Scand J Gastroenterol.* 2006;41(3):302–305. doi:10.1080/00365520500206517.
  100. Casella G, Orfanotti G, Giacomantonio L, et al. Celiac disease and obstetrical-gynecological contribution. *Gastroenterol Hepatol Bed Bench.* 2016;9(4):241–249.
  101. Kotze LM. Gynecologic and obstetric findings related to nutritional status and adherence to a gluten-free diet in Brazilian patients with celiac disease. *J Clin Gastroenterol.* 2004;38:567–574.
  102. Sferlazzas C, Arrigo T, Salzano G, et al. Menarcheal age in celiac disease may not be delayed and may be irrespective of age at diagnosis and dietary management. *J Endocrinol Invest.* 2008;31(5):432–435. doi:10.1007/BF03346387.
  103. Santonicola A, Iovino P, Cappello C, Capone P, Andreozzi P, Ciacci C. From menarche to menopause: the fertile life span of celiac women. *Menopause.* 2011;18(10):1125–1130. doi:10.1097/gme.0b013e3182188421.
  104. Ehsani-Ardakani MJ, Fallahian M, Rostami K, et al. Celiac disease and dysfunctional uterine bleeding; the efficiency of gluten free diet. *Bratislavske lekarske listy.* 2014;115(1):19–21.
  105. Singh P, Arora S, Lal S, Strand TA, Makharia GK. Celiac disease in women with infertility: a meta-analysis. *J Clin Gastroenterol.* 2016;50(1):33–39. doi:10.1097/MCG.0000000000000285.
  106. Moleski SM, Lindenmeyer CC, Veloski JJ, et al. Increased rates of pregnancy complications in women with celiac disease. *Ann gastroenterol.* 2015;28(2):236–240.
  107. Saccone G, Berghella V, Sarno L, et al. Celiac disease and obstetric complications: a systematic review and metaanalysis. *Am J Obstet Gynecol.* 2016;214(2):225–234. doi:10.1016/j.ajog.2015.09.080.
  108. D'Ippolito S, Gasbarrini A, Castellani R, et al. Human leukocyte antigen (HLA) DQ2/DQ8 prevalence in recurrent pregnancy loss women. *Autoimmun Rev.* 2016;15(7):638–643. doi:10.1016/j.autrev.2016.02.009.
  109. Rostami K, Steegers EA, Wong WY, Braat DD, Steegers-Theunissen RP. Coeliac disease and reproductive disorders: a neglected association. *Eur J Obstet Gynecol Reprod Biol.* 2001;96:146–149.
  110. Singhal N, Alam S, Sherwani R, Musarrat J. Serum zinc levels in celiac disease. *Indian Pediatr.* 2008;45:319–321.
  111. Yuce A, Demir H, Temizel IN, Kocak N. Serum carnitine and selenium levels in children with celiac disease. *Indian J Gastroenterol.* 2004;23:87–88.
  112. Cakmak E, Karakus S, Demirpence O, Demet Coskun B. Ovarian reserve assessment in celiac patients of reproductive age. *Med Sci Monitor.* 2018;24:1152–1157.
  113. Di Simone N, De Spirito M, Di Nicuolo F, et al. Potential new mechanisms of placental damage in celiac disease: anti-transglutaminase antibodies impair human endometrial angiogenesis. *Biol Reprod.* 2013;89(4):88. doi:10.1095/biolreprod.112.106542.
  114. Di Simone N, Silano M, Castellani R, et al. Anti-tissue transglutaminase antibodies from celiac patients are responsible for trophoblast damage via apoptosis in vitro. *Am J Gastroenterol.* 2010;105(10):2254–2261. doi:10.1038/ajg.2010.233.
  115. Tersigni C, Castellani R, De Waure C, et al. Celiac disease and reproductive disorders: meta-analysis of epidemiologic associations and potential pathogenic mechanisms. *Hum Reprod Update.* 2014;20(4):582–593. doi:10.1093/humupd/dmu007.
  116. Cooke WT, Peeney AL, Hawkins CF. Symptoms, signs and diagnostic features of idiopathic steatorrhea. *Q J Med.* 1953;22:59–78.
  117. Farthing MJ, Edwards CR, Rees LH, Dawson AM. Male gonadal function in coeliac disease: 1. Sexual dysfunction, infertility, and semen quality. *Gut.* 1982;23:608–614.
  118. Hogen Esch CE, Van Rijssen MJ, Roos A, et al. Screening for unrecognized coeliac disease in subfertile couples. *Scand J Gastroenterol.* 2011;46(12):1423–1428. doi:10.3109/00365521.2011.615858.
  119. Meloni GF, Dessole S, Vargiu N, Tomasi PA, Musumeci S. The prevalence of coeliac disease in infertility. *Human Reprod.* 1999;14:2759–2761.
  120. Khoshbaten M, Rostami Nejad M, Farzady L, Sharifi N, Hashemi SH, Rostami K. Fertility disorder associated with celiac disease in males and females: fact or fiction? *J Obstet Gynaecol Res.* 2011;37(10):1308–1312. doi:10.1111/j.1447-0756.2010.01518.x.
  121. Farthing MJ, Rees LH, Dawson AM. Male gonadal function in coeliac disease: III. Pituitary regulation. *Clin Endocrinol (Oxf).* 1983;19:661–671.
  122. Farthing MJ, Rees LH, Edwards CR, Dawson AM. Male gonadal function in coeliac disease: 2. Sex hormones. *Gut.* 1983;24:127–135.
  123. Green JR, Goble HL, Edwards CR, Dawson AM. Reversible insensitivity to androgens in men with untreated gluten enteropathy. *Lancet.* 1977;1:280–282.
  124. Delvecchio M, Faienza MF, Lonero A, Rutigliano V, Francavilla R, Cavallo L. Prolactin may be increased in newly diagnosed celiac children and adolescents and decreases after 6 months of gluten-free diet. *Horm Res Paediatr.* 2014;81(5):309–313. doi:10.1159/000357064.
  125. Kapur G, Patwari AK, Narayan S, Anand VK. Serum prolactin in celiac disease. *J Trop Pediatr.* 2004;50(1):37–40. doi:10.1093/tropej/50.1.37.

126. Assiri AM. Isolated short stature as a presentation of celiac disease in Saudi children. *Pediatric Reports*. 2010;2(1):e4. doi:10.4081/pr.2010.e4.
127. Bonamico M, Scire G, Mariani P, et al. Short stature as the primary manifestation of monosymptomatic celiac disease. *J Pediatr Gastroenterol Nutr*. 1992;14(1):12–16.
128. Bozzola M, Giovenale D, Bozzola E, et al. Growth hormone deficiency and coeliac disease: an unusual association? *Clin Endocrinol (Oxf)*. 2005;62(3):372–375. doi:10.1111/j.1365-2265.2005.02227.x.
129. Federico G, Favilli T, Cinquanta L, Ughi C, Saggese G. Effect of celiac disease and gluten-free diet on growth hormone-binding protein, insulin-like growth factor-I, and insulin-like growth factor-binding proteins. *Horm Res*. 1997;48(3):108–114. doi:10.1159/000185499.
130. Giovenale D, Meazza C, Cardinale GM, et al. The prevalence of growth hormone deficiency and celiac disease in short children. *Clin Med Res*. 2006;4(3):180–183. doi:10.3121/cmr.4.3.180.
131. Giannattasio A, Did F, Minicucci V, et al. A retrospective evaluation of the association of celiac disease and growth hormone deficiency: more than a casual association? *Minerva Endocrinologica*. 2017;42(1):24–29. doi:10.23736/S0391-1977.16.02366-X.
132. Salardi S, Cacciari E, Volta U, et al. Growth and adult height in atypical coeliac patients, with or without growth hormone deficiency. *J Pediatr Endocrinol Metabol*. 2005;18(8):769–775. doi:10.1515/JPEM.2005.18.8.769.
133. Giovenale D, Meazza C, Cardinale GM, et al. Growth hormone treatment in prepubertal children with celiac disease and growth hormone deficiency. *J Pediatr Gastroenterol Nutr*. 2007;45(4):433–437. doi:10.1097/MPG.0b013e3180de5e0a.
134. Weiss B, Skourikhin Y, Modan-Moses D, Broide E, Fradkin A, Bujanover Y. Is adult height of patients with celiac disease influenced by delayed diagnosis? *Am J Gastroenterol*. 2008;103(7):1770–1774. doi:10.1111/j.1572-0241.2008.01844.x.
135. Meazza C, Pagani S, Messini B, et al. Coeliac children treated for growth hormone deficiency reach normal final height. *Clin Endocrinol (Oxf)*. 2011;74(6):791–792. doi:10.1111/j.1365-2265.2010.03960.x.
136. Ferrante E, Giavoli C, Elli L, et al. Evaluation of GH-IGF-I axis in adult patients with coeliac disease. *Hormone and Metabolic Research = Hormon- Und Stoffwechselforschung = Hormones Et Metabolisme*. 2010;42(1):45–49. doi:10.1055/s-0029-1241169.
137. Delvecchio M, De Bellis A, Francavilla R, et al. Antipituitary antibodies in children with newly diagnosed celiac disease: a novel finding contributing to linear-growth impairment. *Am J Gastroenterol*. 2010;105(3):691–696. doi:10.1038/ajg.2009.642.
138. Husebye ES, Anderson MS, Kampe O. Autoimmune Polyendocrine Syndromes. *N Engl J Med*. 2018;378(12):1132–1141. doi:10.1056/NEJMra1713301.
139. Lakhota M, Pahadia HR, Kumar H, Singh J, Tak SA. Case of Autoimmune Polyglandular Syndrome (APS) Type II with hypothyroidism, hypoadrenalism, and celiac disease - a rare combination. *J Clin Diagn Res*. 2015;9(4):OD01–3. doi:10.7860/JCDR/2015/10755.5748.
140. Hrubisková K, Jackuliak P, Vanuga P, Pura M, Payer J. Autoimmune polyendocrine syndrome type 2 associated with autoimmune hypophysitis and coeliac disease. *Vnitr Lek*. 2010;56:1169–1176.
141. Kirmızibekmez H, Yeşiltepe Mutlu RG, Demirkıran Urgancı N, Öner A. Autoimmune polyglandular syndrome type 2: a rare condition in childhood. *J Clin Res Pediatr Endocrinol*. 2015;7(1):80-2. doi:10.4274/jcrpe.1394.
142. Maturu A, Michels A, Draznin B. Multiple disease associations in autoimmune polyglandular syndrome type II. *Endocr Pract*. 2014;20(12):e250–5. doi:10.4158/EP14159.CR.
143. Dieli-Crimi R, Núñez C, Estrada L, López-Palacios N. An autoimmune polyglandular syndrome complicated with celiac disease and autoimmune hepatitis. *Ann Hepatol*. 2016;15(4):588–91.
144. Innico G, Frasseti N, Coppola B, Mariotti A, Lai S. Autoimmune polyglandular syndrome in a woman of 51 years. *Eur Rev Med Pharmacol Sci*. 2014;18:1717–1719.
145. Noble JA, Valdes AM, Cook M, Klitz W, Thomson G, Erlich HA. The role of HLA class II genes in insulin-dependent diabetes mellitus: molecular analysis of 180 Caucasian, multiplex families. *Am J Hum Genet*. 1996;59:1134–1148.
146. Simmonds MJ, Gough SC. Unravelling the genetic complexity of autoimmune thyroid disease: HLA, CTLA-4 and beyond. *Clin Exp Immunol*. 2004;136(1):1–10. doi:10.1111/j.1365-2249.2004.02424.x.
147. Erichsen MM, Lovas K, Skiningsrud B, et al. Clinical, immunological, and genetic features of autoimmune primary adrenal insufficiency: observations from a Norwegian registry. *J Clin Endocrinol Metab*. 2009;94(12):4882–4890. doi:10.1210/jc.2009-1368.
148. Zhao Z, Zou J, Zhao L, et al. Celiac disease autoimmunity in patients with autoimmune diabetes and thyroid disease among chinese population. *PLoS One*. 2016;11(7):e0157510. doi:10.1371/journal.pone.0157510.
149. Kurien M, Mollazadegan K, Sanders DS, Ludvigsson JF. Celiac disease increases risk of thyroid disease in patients with type 1 diabetes: a nationwide cohort study. *Diabetes Care*. 2016;39(3):371–375. doi:10.2337/dc15-2117.
150. Saha S, Saini S, Makharia GK, Datta Gupta S, Goswami R. Prevalence of coeliac disease in idiopathic hypoparathyroidism and effect of gluten-free diet on calcaemic control. *Clin Endocrinol (Oxf)*. 2016;84(4):578–586. doi:10.1111/cen.12850.
151. Somani S, Kotwal N, Upreti V. An unusual case of hypercalcemia in a patient of concomitant hypoparathyroidism and celiac disease. *Clin Cases Min Bone Metabol*. 2016;13(1):51–53. doi:10.11138/ccmbm/2016.13.1.051.

152. Yerushalmi B, Lev-Tzion R, Loewenthal N. Refractory hypoparathyroidism in a child with celiac disease. *Israel Med Assoc J.* 2016;18:58–60.
153. Greenfield JR, Samaras K. Evaluation of pituitary function in the fatigued patient: a review of 59 cases. *Eur J Endocrinol.* 2006;154(1):147–157. doi:10.1530/eje.1.02010.
154. Collin P, Hakanen M, Salmi J, Maki M, Kaukinen K. Autoimmune hypopituitarism in patients with coeliac disease: symptoms confusingly similar. *Scand J Gastroenterol.* 2001;36:558–560.
155. Kumar V, Valeski JE, Wortsman J. Celiac disease and hypoparathyroidism: cross-reaction of endomysial antibodies with parathyroid tissue. *Clin Diagn Lab Immunol.* 1996;3:143–146.
156. Lee RH, Lyles KW, Colon-Emeric C. A review of the effect of anticonvulsant medications on bone mineral density and fracture risk. *Am J Geriatr Pharmacother.* 2010;8(1):34–46. doi:10.1016/j.amjopharm.2010.02.003.
157. Verrotti A, Loiacono G, Laus M, Coppola G, Chiarelli F, Tiboni GM. Hormonal and reproductive disturbances in epileptic male patients: emerging issues. *Reprod toxicol.* 2011;31(4):519–527. doi:10.1016/j.reprotox.2011.02.002.
158. Aggarwal S, Lebwohl B, Green PH. Screening for celiac disease in average-risk and high-risk populations. *Therap Adv Gastroenterol.* 2012;5(1):37–47. doi:10.1177/1756283X11417038.